

Cambridge South East Transport Phase 2 Environmental Statement

Appendix 10.12 Biodiversity Net Gain Assessment 31st July 2023



Methodology

Desk study

10.10.1 The following resources were reviewed to inform the assessment of the baseline habitat types and condition assessment using the Biodiversity Metric 3.1:

- Aerial imagery and OS maps;
- Multi-Agency Geographic Information for the Countryside (MAGIC);
- Environment Agency Catchment Data Explorer,
- 10.10.2 For areas of the Site where BNG-specific field surveys were not undertaken a desk-based conversion of Phase 1 Habitat codes (determined through the Phase 1 Habitat survey carried out between 2018 and 2020) and/ or aerial imagery to a metric specific habitat type and condition was undertaken.
- In relation to this BNG assessment, the desk study undertaken in March 2023 was used to 10.10.3 evaluate the potential for important ecological features, and how their presence may influence the feasibility of delivering BNG, within and adjacent to the Site. Impacts to statutory designated sites for nature conservation and irreplaceable habitats are not adequately measured by Biodiversity Metric 3.1 and they require separate consideration that complies with relevant policy and legislation.
- 10.10.4 To identify any designated sites for nature conservation within 2 km of the Site and irreplaceable habitats and/ or priority habitats within the Site, the following online data sources were accessed:
 - MAGIC¹
 - Ancient Tree Inventory²
- 10.10.5 The Metric applies a strategic significance score based on whether the location of a site and/ or the baseline habitats present, or proposed habitat interventions have been identified as being locally important for biodiversity. All habitat parcels must be assigned a strategic significance score. The options for scoring each habitat parcel are:
 - High formally identified in local strategy, plan or policy (or River Basin Management) Plan, Catchment Plan and/or Local Plans);
 - Medium location ecologically desirable but not identified in a local strategy, plan or policy; or
 - Low not identified in a local strategy, plan or policy OR no strategy or plan is in place in the area.
- 10.10.6 The desk study was used to identify local biodiversity strategies and plans, as set out by Cambridge City Council and Greater Cambridge Shared Planning to determine the strategic significance of the Site and/ or habitats within the Site. The following online data sources were accessed:
 - Cambridge City Council Local Plan (2018)
 - South Cambridgeshire Local Plan (2018)
 - Greater Cambridge Local Plan First Proposals (2021)
 - Greater Cambridge Local Plan Development Strategy Update report (2023) •
 - Greater Cambridge Biodiversity Supplementary Planning Document •

- Cambridge City Council Biodiversity Strategy 2022-2030
- Buglife's B-Lines

Area habitat and linear hedgerow survey and condition assessment

10.10.7 The baseline biodiversity habitat data for the Site was calculated using habitat mapping and condition assessment data between September and November 2022. The condition assessment survey involved mapping habitats present on Site using a modified UK Habitat (UKHab) Classification System as set out within the Biodiversity Metric 3.1 technical supplement and assigning a condition according to the methodology and criteria outlined within this technical supplement.

10.10.8 As part of the ecological walkover survey, the 'condition', as set out by the Metric, of each habitat, was recorded for each area habitat and linear hedgerow habitat. The habitat type was assessed and categorised as either 'Good', 'Fairly Good', 'Moderate', 'Fairly Poor', or 'Poor'. The assessment was made using the criteria within the Metric's Technical Supplement³, with specific assessment criteria for each broad habitat type.

- 10.10.9 If a habitat type varied in condition within the Site, this was recorded and mapped separately. The area covered by each habitat type was mapped using Geographical Information Systems (GIS).
- 10.10.10 The Minimum Mappable Unit (MMU) was 25 m² for area habitats and 5 m length x 1 m width for linear features. Secondary codes were utilised focussing on habitat features and land use.
- 10.10.11 A UKHab plan of the Site developed for the Metric, which includes hedgerows assessed as linear features is provided in Annex A.

Modular River Physical (MoRPh) survey

- 10.10.12 The baseline condition of the River Granta was surveyed by WSP using the MoRPh survey method as directed by the Biodiversity Metric guidance. The River Granta was divided into discrete reaches for the purpose of MoRPh survey siting and metric reporting, taking account of the Site interactions and existing river typology.
- MoRPh surveys were conducted in September and October 2022 by an accredited 10.10.13 MoRPh surveyor. The MoRPh method is a quantitative visual geomorphological assessment of a river and its riparian zone (up to 10 m from the bank edge), that records a list of features which are given categories (as follows) based on their extent across the survey sub-reach:
 - Extensive (>33%);
 - Present (5% to 33%): •
 - Trace (<5%);
 - Absent. •
- 10.10.14 Such features include elements such as channel form, in-channel habitats (e.g. riffles and pools), bed substrates and bank material, as well as flow types. Broad categories of aquatic plant community structure and characteristics of the bankside and riparian zone are also recorded.

2 Woodland Trust (2023) Ancient Tree Inventory [online] Available at: https://ati.woodlandtrust.org.uk/ 3 Buglife [online] Available at: https://statics.teams.cdn.office.net/evergreen-assets/safelinks/1/atp-safelinks.html

¹ Multi-Agency Geographic Information for the Countryside (MAGIC) [Online]. Available at: https://magic.defra.gov.uk/

10.10.15 MoRPh surveys were undertaken at or adjacent to Proposed Development interactions, in this case at Proposed Development crossing locations. Three MoRPh surveys were conducted on the River Granta at/ near to the Babraham crossing (Reach D1), and two MoRPh surveys were undertaken on the River Granta at/ near to the Stapleford crossing (Reach D2), with one of the MoRPh surveys in each location coincident with the crossing location (D1-S3 and D2-S2). Reaches and sub-reaches have been numbered from upstream to downstream. A minimum of 20% of the river length was covered within the Red Line Boundary (RLB) limits. Each sub-reach survey was 50 m in length and comprised of five contiguous MoRPh modules of 10 m. The survey locations and subreach codes are presented in Table A10.12.1

Watercourse	Reach	Sub- reach	NGR	Date Surveyed
River Granta	D1	S1	TL 51819 49346 to TL 51788 49392	29/09/2022
		S2	TL 51674 49627 to TL 51624 49625	06/10/2022
		S3	TL 51374 49673 to TL 51326 49696	06/10/2022
	D2	S1	TL 48780 51289 to TL 48729 51282	29/09/2022 & 04/10/2022
		S2	TL 48438 51236 to TL 48390 51231	04/10/2022

Table A10.12.1 MoRPh survey details

10.10.16 The River Granta adjoins the Site along two sections: the sub-reaches D1-S1 and D2-S1. The Biodiversity Metric guidance states that:

> "Where the red line boundary of the development encompasses the riparian zone, either whole or in part, but excludes the channel of the watercourse, the rivers and streams metric (including the condition assessment) must be applied."

- 10.10.17 Therefore, River Biodiversity Units (RBUs) have been calculated for these reaches and they have been treated as being within the Site. Note that this guidance is not relevant to ditches as the riparian zone of a ditch is not used to influence its condition score, therefore any ditches which adjoin but are outside of the Site were not included.
- 10.10.18 Note that due to a revision in the Site boundary that occurred after the surveys were completed, one of the MoRPh sub-reaches (D1-S2) is outside of the current RLB. The results of this survey are presented to provide contextual information on the condition of the wider River Granta but have not been used to calculate RBUs.
- 10.10.19 The final river condition scores for rivers within the RLB limits were inputted into the Natural England Biodiversity Metric 3.1.

Ditch condition assessment

- 10.10.20 The baseline condition of watercourses identified as ditches were surveyed using the ditch condition assessment as directed by the Biodiversity Metric guidance.
- The ditch condition assessment considers the following criteria in assigning condition: 10.10.21
 - The ditch is of good water guality, with clear water (low turbidity) indicating no obvious signs of pollution;

- A range of emergent, submerged and floating leaved plants are present. As a guide, >10 species of emergent, floating or submerged plants in a 20 m ditch length;
- There is less than 10% cover of filamentous algae and/ or duckweed (these are signs of eutrophication);
- A fringe of marginal vegetation is present along more than 75% of the ditch;
- Physical damage evident along less than 5% of the ditch, such as excessive poaching, damage from machinery use or storage, or any other damaging management activities;
- Sufficient water levels are maintained; as a guide a minimum summer depth of approximately 50 cm in minor ditches and 1 m in main drains;
- Less than 10% of the ditch is heavily shaded; •
- There is absence of non-native plant and animal species . •
- 10.10.22 All eight criteria must be passed in order to achieve Good condition. If a ditch passes six or seven of the criteria it achieves Moderate condition. Any ditch that passes five or less of the criteria has a condition class of Poor. The survey locations and sub-reach codes are presented in Table A10.12.2.

Table A10.12.2 Locations and sub-reach codes of the watercourses assessed

Watercourse	Sub-reach	Mid-point NGR	Dry at time of survey?*
Hobson's Brook	D4	TL 45994 54160	No
Unnamed ditches	D2E	TL 48548 51143	Yes
	D5A	TL 46377 53409	NA*
	W1	TL 45961 55035	No
	W2	TL 45982 54398	Yes
	W3	TL 48025 51401	Yes
	W4	TL 48667 50898	No
	W5	TL 48840 50703	Yes
	W6	TL 48323 51353	Yes
	W7	TL 45914 54402	NA*
	W8	TL 46052 54487	No
	W9	TL 45969 55029	No
	W22	TL 46413 53296	NA*
	W24	TL 46004 54883	Yes
	W27	TL 46113 54195	NA*

Watercourse	Sub-reach	Mid-point NGR	Dry at time of survey?*
	W29	TL 46081 54623	No

*These ditches were not surveyed due to access constraints or because surveys were conducted prior to a change of Site boundary

- 10.10.23 Hobson's Brook was surveyed using the ditch condition assessment method due to the watercourse presenting a ditch-like typology at the location of its interaction with the Proposed Development. However, a review of aerial imagery and historical mapping indicates that the watercourse is naturally present and has been existing in the landscape as far back as the 1885-1900 map series. Consequently, it does not meet the definition of ditch according to the Biodiversity Metric guidance. Hobson's Brook must, therefore, be considered within the rivers and streams broad habitat typology, for which purpose ditch condition assessment does not provide an appropriate condition score. The condition class of Hobson's Brook has therefore been assigned based upon professional judgement and review of available data.
- 10.10.24 Several of the ditches were dry at the time of survey but have been given a precautionary condition assessment due to the unusually dry conditions in the year preceding the survey, i.e., it is assumed that surveys were undertaken during an atypical year and that during a typical year the ditches are likely to retain water for more than four months of the year4. This may result in an overestimation of baseline RBUs for ditches.
- 10.10.25 Ditches 5A and W7 were not possible to survey due to access limitations. Similarly, ditches W22 and W27 were outside of the RLB at the time of survey and were therefore excluded from field survey, however, these ditches are now within the RLB. Ditch condition has therefore been applied to these ditches based upon the condition of nearby ditches, aerial imagery, information from previous water vole and otter surveys carried out in 2022, and professional judgement. It was determined that ditches were of a similar character and condition to the other ditches within the RLB.
- 10.10.26 The final ditch condition scores for ditches within the RLB limits were inputted into the Natural England Biodiversity Metric 3.1.

On-site post development habitats

- 10.10.27 The current general arrangement and landscape design plans and management plans have been used to inform the post development scenario:
 - Cambridge South East Transport Phase 2 TWAO Application Public Transport and Active Travel Corridor Bridges, Structures and Proposed Development Extents5
 - Cambridge South East Transport Phase 2 TWAO Application Public Transport and Active Travel Corridor – Route Layout Sheet 1 to 186
 - Drainage Retention Pond drawing7
 - Flood Compensation drawing8
 - Landscape and Environmental Management Plan (ES Volume 2, Appendix 2.1)

10.10.28 These plans were converted to a GIS environment and combined to produce a single layer, ensuring gaps and overlaps were removed, where possible. From this layer the areas of retained habitats, new carriageway and temporary land take were calculated. Each proposed planting type was translated into a UKHab classification category using the professional judgement of Atkins Ecologists and a target condition assigned based on the likely achievable condition, as outlined in Table A10.12.3. The likely achievable condition was determined with reference to the criteria outlined within the Biodiversity Metric 3.1 technical supplement. Areas of proposed post development intervention (habitat creation and/ or habitat retention/enhancement), including built development, were calculated using GIS.

Table A10.11.3 Translation of the landscape schedule to UKHabs metric

Landscape planting type	Metric habitat	Assumed co		
Area habitat				
Chalk grassland	Lowland calcareous grassland	Moderate – A grass and wil as high-quali moderate tar		
Amenity grassland	Modified grassland	Poor - Althou of species it v and has beer precaution.		
Species-rich grassland	Other neutral grassland	Moderate – s native floweri is likely to be areas were p non-cereal cr any agricultur years followir proposed roa be subject to this habitat ty condition/price		

5 Drawing Number 5212868-ATK-GEN-WHL AL SCHME-DR-CX-000001. Rev P01.1 6 Drawing Number 5212868-ATK-HGN-WHL AL SCHME-DR-CH-000001 to 000018. Rev C01 7 Drawing Ref CD-Ss 50 70 95 DGRP-M-Dn DrainageRetentionPond

ondition and justification

Although this will have a diverse range of Idflower species it is unlikely to qualify ity priority grassland and has been given get condition as a precaution.

ugh this is likely to have a good number will be subject to regular management n given moderate target condition as a

species rich grassland mix will include ing plants. Although perennial rye grass present at less than 30% a lot of the previously cropland including cereal and rops therefore existing seedbank and ral weeds likely to germinate in first ng planting. This habitat type includes adside planting and are therefore likely to disturbance and management. As such ype is unlikely to achieve good ority habitat status.

⁴ Ditches are defined as - 'Artificially created, linear water-conveyancing features that are less than 5 m wide and likely to retain water for more than 4 months of the year. Their hydraulic function is primarily for land drainage, and although partially or fully connected to a river system, they would not have been present without human intervention'. Definition after Panks et al. (2022)

⁸ Drawing Reference ATK-HWY_Flood Compensation

Landscape planting type	Metric habitat	Assumed condition and justification	
Ponds	Ponds (non- priority habitat)	Moderate – By the nature of the inclusion in the design as drainage retention or flood compensation, the ponds are likely to be subject to pollution and run-off from the road and therefore given a precautionary moderate condition. Other criteria are expected to be met, including natural edge habitat, less than 10% duckweed or filamentous algae, absence of invasive species, water levels allowed to fluctuate and not artificially connected to other waterbodies or stocked with fish.	
Woodland / Field margin	Other woodland; broadleaved	Moderate - Plantation woodland that is unlikely to have a diverse age and height structure or presence of standing and fallen deadwood so unlikely to me good condition. Expected to meet other condition assessment criteria, including dominance of native species, complete canopy cover, free from damage or inappropriate management, and absence of invasive species.	
Hedgerow	Modified grassland / Other neutral grassland	Moderate – Hedgerow habitats were mapped as hedgerow area habitats within the landscape plan. Therefore, this has been converted to grassland area habitat with hedgerow features mapped on top as linear hedgerow (outlined separately below), in line with the Biodiversity Metric 3.1 User Guide.	
Linear hedgerow habitat			
N/A	Native species-rich hedgerow	Moderate – Hedgerows largely comprise the boundary features along the Proposed Development and adjacent land. Therefore, subject to regular management and possible nutrient enrichment from farmland/roads as such unlikely to meet criteria for good condition and also unlikely to qualify as a priority habitat.	

- 10.10.29 It is assumed that any land lying outside of these areas (i.e. within the temporary land take) would be returned to landowners in the same condition as the baseline.
- 10.10.30 The above general arrangement and landscape design plans and management plans were also used to identify points of interaction between the Proposed Development and linear river features.

Calculating area and linear hedgerow biodiversity units using Biodiversity Metric 3.1

- 10.10.31 Biodiversity Metric 3.1 uses a number of measures to quantify baseline biodiversity value for each habitat type within a site boundary. These measures include the habitats' intrinsic value (i.e. its distinctiveness), its condition and its area/ length (ha / km). In addition, the local strategic significance of a site and/or the habitats within it is taken into account by applying a spatial multiplier.
- 10.10.32 Area-based habitat features, linear hedgerow features (which includes lines of trees) and linear river features (which includes rivers, streams, ditches and canals) are measured as separate components in the Metric and the delivery of units, including the overall net unit change for a site, are reported separately.
- 10.10.33 A habitat's distinctiveness score is derived from its intrinsic biodiversity value, reflecting the rarity of the plant community, the time it takes to reach maturity, its value to fauna, and its ecosystem function. This score is pre-assigned and is pre-populated in the Metric.
- 10.10.34 The calculation of baseline biodiversity units for area-based habitat and linear hedgerow features is shown in Figure A10.12.1. The calculation for linear river features includes additional multipliers that take into account whether a river feature has been modified by built development through the presence of encroachment into the watercourse channel or riparian zone, i.e. the multipliers reduce the number of river units to reflect the negative impact that such encroachment has on a naturally functioning river system.



Figure A10.12.1 Calculation of Biodiversity Units using Biodiversity Metric 3.1. NB. Illustration does not fully depict calculation of river units; additional multipliers taking into account watercourse and riparian encroachment are applied. Text underlined indicates a risk factor applied when creating or enhancing habitats. *Applies to off-site habitat interventions only.

Post development biodiversity units are calculated the same way but with the addition of 10.10.35 risk multipliers that take into account the difficulty and time it takes to create new habitat or enhance existing habitat (see Figure A10.12.1). These temporal and risk related multipliers are set by Natural England within the Metric and cannot be changed (full details are provided in the Technical Supplement of Biodiversity Metric 3.1). Where habitat creation or enhancement is off-site, biodiversity units are calculated the same way but a spatial risk multiplier is applied based on the distance of habitat creation or enhancement to the site of habitat loss. No off-site habitat creation or enhancement is proposed for the Proposed Development.

- 10.10.36 The Metric also takes into account whether habitat creation or enhancement is delivered in advance of any impact it is compensating for or whether there will be any significant delay in an intervention relative to the impact. If taken into account, the standard temporal risk multiplier is adjusted accordingly.
- 10.10.37 To calculate the overall net change in biodiversity units, the baseline units are subtracted from the post development units, as detailed in Figure A10.12.1.

Calculating baseline river biodiversity units using the Biodiversity Metric 3.1

- 10.10.38 The Biodiversity Metric 3.1 includes a Rivers and Streams Module³⁰. As for terrestrial habitats, the Rivers and Streams Module requires an assessment of baseline River Biodiversity Units (RBU) and post-works RBUs in order to assess overall change in biodiversity units.
- To determine the baseline RBUs, four key components are assessed and inputted to the 10.10.39 Natural England Biodiversity Metric 3.1 Calculation Tool (along with the length of rivers/ streams and ditches within the RLB). These scores result in weightings which are used to calculate the number of RBUs. The components are:
 - Distinctiveness
 - Strategic significance •
 - Watercourse and riparian encroachment •
 - River/ ditch condition
- 10.10.40 The distinctiveness score is based on the type of river habitat present, and its value based on its rarity. In the Rivers and Streams Module of the Metric, rivers can be classed as 'very high', 'high', 'medium' or 'low'. All ditches have a default classification of low distinctiveness.
- 10.10.41 Strategic significance is dependent on whether the watercourse has identified actions within River Basin Management Plans (i.e. is a delineated Water Framework Directive (WFD) reportable water body), Catchment Plans and/ or Local Plans, meaning that the watercourse has 'High strategic significance'. If this is not the case, the watercourse is considered to have 'Low strategic significance'.
- 10.10.42 Encroachment scores refer to the extent of encroachment of a development: (a) in the watercourse, and (b) in the riparian zone (10 m from the top of the riverbank). Encroachment within the watercourse is classified as 'no encroachment', 'minor' or 'major' whereas in the riparian zone encroachment is classed as 'no encroachment', 'minor', 'moderate' or 'major'.
- 10.10.43 River condition is based upon results from MoRPh surveys. Data collected during surveys were uploaded to the MoRPh cartographer workspace. Cartographer calculates 32 indicators of river condition for each MoRPh sub-reach which are scaled according to river type (determined through the desk study component of the river condition assessment), to produce the final river condition score for each sub-reach. The final river condition score produced in the assessment may be: 'Good', 'Fairly Good', 'Moderate', 'Fairly Poor' or 'Poor'. Ditch condition is based upon ditch condition assessments, with possible scores of 'Good', 'Moderate' or 'Poor' assigned based upon the number of criteria met.
- 10.10.44 Further details on the methods for determining distinctiveness, encroachment, strategic significance and condition are detailed in the Biodiversity Metric 3.1 guidance.

- 10.10.45 RBUs are linear units and cannot be combined with the area units calculated for terrestrial habitats within the assessment. A net gain in RBUs would not automatically correlate to an overall net gain for the project as there may be a net loss in terrestrial biodiversity units. Therefore, each of the different module components of the Metric need to be considered in parallel and reported separately.
- 10.10.46 This is also true for rivers/ streams and ditches, as they have different distinctiveness. As per the Trading Rules under the Biodiversity Metric 3.1 guidance. RBUs cannot be traded across river types of different distinctiveness. As such, rivers/ streams and ditches have been assessed separately within this iteration of the BNG assessment.

Mitigation hierarchy

- In accordance with good practice, the principles of the mitigation hierarchy and the 10.10.47 avoidance of impacts to irreplaceable habitats must be applied when committing to achieving BNG. This is reinforced through the publication of the British Standard BS 8683:20211.
- 10.10.48 The sequential approach to applying the mitigation hierarchy is to:
 - Avoid Where possible habitat damage should be avoided;
 - Minimise Where possible habitat damage and loss should be minimised;
 - Remediate Where possible any damaged or lost habitat should be restored; •
 - Compensate As a last resort, damaged or lost habitat should be compensated for .
- Application of the mitigation hierarchy is encouraged by the Metric because BNG can be 10.10.49 more easily achieved by avoiding the loss of habitat through either retention or enhancement.

BNG Good Practice Principles

10.10.50 To claim BNG, evidence that demonstrates a project has clearly applied all ten of the good practice principles for achieving BNG, as detailed in Table A10.12.4, must be provided. Adhering to all principles together underpins the good practice required for achieving and sustaining BNG.

Table A10.12.4 BNG good practice principles

Principle	In practice
Apply the mitigation hierarchy	Do everything possible to first avoid and then minimise impacts on biodiversity. Only as a last resort, and in agreement with external decision-makers where possible, compensate for losses that cannot be avoided. If compensating for losses within the development footprint is not possible or does not generate the most benefits for nature conservation, then offset biodiversity losses by gains elsewhere.
Avoid losing biodiversity that cannot be offset by gains elsewhere	Avoid impacts on irreplaceable biodiversity – these impacts cannot be offset to achieve no net loss or net gain.
Be inclusive and equitable	Engage stakeholders early, and involve them in designing, implementing, monitoring and evaluating the approach to net gain. Achieve net gain in partnership with stakeholders where possible and share the benefits fairly among stakeholders.
Address risks	Mitigate difficulty, uncertainty and other risks to achieving net gain. Apply well-accepted ways to add contingency when calculating biodiversity losses and gains in order to account for any remaining risks, as well as to compensate for the time between losses occurring and the gains being fully realised.
Make a measurable net gain contribution	Achieve a measurable, overall gain for biodiversity and the services ecosystems provide while directly contributing towards nature conservation priorities.

Principle	In practice
Achieve the best outcomes for biodiversity	Achieve the best outcome credible evidence and loc choices when:
	Delivering compensation amount and condition and timing of biodiversity loss
	Compensating for losses a different type that delive conservation;
	Achieving net gain locally contributing towards nature regional and national lever
	Enhancing existing or cre
	Enhancing ecological con better and joined areas fo
Be additional	Achieve nature conservat exceed existing obligation would occur anyway).
Create a net gain	Ensure net gain generate
legacy	Engaging stakeholders ar that secure net gain in pe
	Planning for adaptive man funding for long-term man
	Designing net gain for bio factors, especially climate
	Mitigating risks from other
	Avoiding displacing harms another;
	Supporting local-level ma
Optimise sustainability	Prioritising BNG and, whe environmental benefits for
Be transparent	Communicate all net gain manner, sharing the learn

9 For development in the UK, the expectation is that compensation sites will be secured for at least the lifetime of the development (e.g. often 25-30 years) with the objective of net gain management continuing in the future.

les for biodiversity by using robust, cal knowledge to make clearly justified

a that is ecologically equivalent in type, and that accounts for the location and ses;

s of one type of biodiversity by providing ers greater benefits for nature

y to the development while also ure conservation priorities at local, els;

eating new habitat;

nnectivity by creating more, bigger, or biodiversity.

ation outcomes that demonstrably ons (i.e. do not deliver something that

es long-term benefits by:

and jointly agreeing practical solutions erpetuity9;

anagement and securing dedicating inagement;

odiversity to be resilient to external e change;

er land uses;

nful activities from one location to

anagement of net gain activities.

ere possible, optimise the wider or a sustainable society and economy.

n activities in a transparent and timely ning with all stakeholders.

Assumptions and Limitations

- 10.10.51 The optimum survey period for habitats is April to October, inclusive. Although habitat type and condition assessments were carried out late September to early October, within the optimum period, some plant species may not be evident due to dormancy. A survey visit was carried out in November, outside the optimum season, in the north of the Site. However, as this is a predominantly urban area with small areas of immature plantation, this is not considered to be a significant constraint.
- 10.10.52 Not all habitat areas received a field survey to assess the habitat types and condition due to a change in the Site boundary following field surveys in September and October 2022. Where this is the case, a desk-based assessment was completed, and a precautionary approach taken to the condition. As such this is not considered to be a significant limitation.
- 10.10.53 Criteria assessment sheets for baseline habitat conditions were not available at the time of writing this BNG assessment and therefore the justification for baseline condition on Site cannot be confirmed. The baseline condition scores have been assessed using professional judgement and reported in the Habitat Type and Condition Assessments for Biodiversity Net Gain Calculations report in line with the Metric Technical Supplement. These baseline conditions were obtained from field and desk-based assessment (where field surveys did not cover the full extent due to a change in site boundary, as described above) and were reflected in the Metric for this BNG assessment, therefore this is not considered to be a significant limitation.
- 10.10.54 The temporary land take will be over two years, so these habitats have been recorded as being lost and subsequently created to reflect the baseline. Therefore, this is not considered to be a significant limitation and has been accounted for in the calculations.
- 10.10.55 Landscape plans depicted hedgerows as area-based habitats. These have been converted to grassland with hedgerows plotted as a linear feature on top. The Metric includes separate calculations for area habitats and linear hedgerows to obtain a more meaningful measure of the ecological value of linear habitats as corridors. Therefore, this is not a significant limitation and has been accounted for in the calculations.
- 10.10.56 For this assessment, it is assumed that habitat creation will be delayed by two years as a worst case scenario (the duration of the anticipated construction programme) from the time of the initial Site clearance and the standard time to target condition has been adjusted accordingly. As detailed design progresses, opportunities to phase the Site clearance and habitat creation proposals will be sought to maximise the predicted delivery of biodiversity units. Based on this, the GCP target net gain of 20% for area habitat and hedgerow habitats is still achievable, however, any project design change will be reflected in the calculation of biodiversity units accordingly.
- It is assumed that an appropriate BNG management and monitoring plan will be 10.10.57 implemented for the duration of the maximum time for the habitat creation proposals to reach their target condition (i.e. 30 years for proposed 'Lowland mixed deciduous woodland' creation) and to give surety that the project will deliver the predicted net gain in biodiversity units. It is then assumed that the habitats will be subject to suitable management to maintain that condition for the operational life of the Proposed Development. A Landscape and Ecological Management Plan (LEMP) (ES Volume 3, Appendix 2.1) has been produced with details of the management to be implemented.
- Weather conditions in 2022 included a prolonged drought meaning that water levels were 10.10.58 particularly low in watercourses. MoRPh scores and ditch condition assessments may be lower than typical as a result.

- 10.10.59 Many of the ditches were dry at time of survey but were still given a precautionary habitat and condition assessment as they were assessed as likely to retain water for more than four months of the year during a typical year, taking into account the antecedent dry conditions. Additionally, the assessment of some ditches was based upon previous surveys or site information as they were not visited during surveys, either due to a change in the RLB or because site access was not possible.
- 10.10.60 The condition of Hobson's Brook has been assigned based upon the ditch condition assessment, aerial imagery, and the WFD status of the water body, because a MoRPh survey was not completed at the point where the watercourse crosses the RLB.

Results

Designated sites

10.10.61 The desk study identified six statutory designated sites for nature conservation within 2 km of the Site. Of these, four were Sites of Special Scientific Interest (SSSI) and two were Local Nature Reserves (LNR). These are detailed in Table A10.12.5.

Table A10.12.5 Statutory designated sites for nature conservation within 2 km of the Site

Site name	Designation	Approximate Distance from Proposed Development	Description
Nine Wells	Local Nature Reserve (LNR)	Adjacent to the RLB	Nine Wells is a historically important site containing several chalk springs, which form the source of the Hobson Conduit. Nine Wells once contained some rare freshwater invertebrates, however following the drought of 1976 these were lost. Today the chalk watercourses are being managed with the aim of re-creating the conditions favourable for a possible re-introduction of these rare species.

site name	Designation	Approximate Distance from Proposed Development	Description	Site name	Designation	
wston Hall eadows	Site of Special Scientific Interest (SSSI)	0.96 km west	This area of meadows overlying spring-fed peat over chalk is a relic of a once common type of wet pasture land found at the margins of the East Anglian Fenland. The plant and animal communities which it supports are generally scarce and poorly represented in Cambridgeshire.	Dernford Fen	SSSI	1
			Grassland communities range from marshy grassland to drier calcareous grassland.			
			The site is also noted for presence of nationally rare umbellifer <i>Selinum</i> <i>carnifolia</i> , now only found in Cambridgeshire. Also of note is the presence of saw wort <i>Serratula</i> <i>tinctoria</i> and the great fen-sedge <i>Cladium mariscus</i> .			
Gog Magog Golf Club	SSSI	1.06 km east	This site supports grassland communities of the calcareous chalk grassland type. Management of the site has secured species-rich calcareous grassland on the Gog Magog Hills overlying chalk and chalky loam over glacial drift.	The Beechwoods	LNR	
			Of additional note is the occurrence of the nationally rare moon carrot <i>Seseli libanotis</i> and the locally rare perennial flax <i>Linum anglicum</i> .			

Description

This site represents a relic of a much larger area of rough fen and carr. Vegetation ranges from dry grassland and scrub to relic fen, reedbed and alder carr. The grassland is of the neutral type on calcareous loam.

The relic fen areas hold reed *Phragmites communis*, saw sedge Cladium mariscus and marsh orchids *Dactylorhiza* spp. Of additional note is the presence of two uncommon fenland orchids, the marsh helleborine *Epipactis palustris* and the fragrant orchid *Gymnadenia conopsea* subsp. *densiflora*.

Areas of open pools within the site together with ditches and the chalk stream along the boundary further enhance the diversity of this site.

A small wood of majestic beeches on a chalk ridge above Cambridge which was planted in the 1840s. Local people planted a new plantation of broad leaved trees in 1992. Wildlife includes white helleborine orchid, beech trees, fungi, great spotted and green woodpeckers, nuthatch, spotted flycatcher. In good beech-mast crop years large flocks of bramblings gather to eat the beech-mast.

Site name	Designation	Approximate Distance from Proposed Development	Description
Roman Road	SSSI	1.8 km east	The Roman Road SSSI supports species-rich calcareous grassland communities of a type which was once widespread on the chalk areas of lowland England and which is now scarce due to changes away from the traditional sheep grazing economy of these areas to arable. Thick hedgerows and small copses along this 'green lane' enhance the value of the grassland for invertebrates.

10.10.62 A total of 24 non-statutory designated sites for nature conservation were identified within 2 km of the Site as detailed in Table A10.12.6.

Table A10.12.6 Non-statutory	designated	sites fo	or nature	conservation	within	2 km
of the Site	-					

Site name	Designation	Approximate distance from Site	Description
River Granta	County Wildlife Site (CWS)	Within the RLB	Major river not grossly modified by pollution or canalisation. Additionally, it supports concentrations of mature pollarded willows <i>Salix</i> sp.
Shelford – Haverhill disused railway (Pampisford)	CWS	155 m to the south-west of the Site, south of Babraham	Supports populations of a Nationally Rare vascular plant <i>Filago</i> <i>pyramidata</i> , and supports frequent numbers of at least six strong calcareous grassland indicator species .
Triangle North of Long Road	CWS	550 m north	The site qualifies as a CWS because of the presence of a Nationally Scarce vascular plant species (<i>Torilis arvensis</i>).
Stapleford Chalk Pit	CWS	750 m to the east	The pit is a site of significance in the wider landscape as it is one of many former chalk quarries in villages across the county, that are now a network of chalk grassland fragments.

Site name	Designation	Approximate distance from Site
Magog Downs Chalk Grassland	CWS	1.0 km to the east
Netherhall Farm Meadow	CWS	1.3 km east
Magog Down and Stapleford Pit	CWS	1.3 km north
Dernford Fen Grassland	CWS	1.46 km east
Wandlebury	CWS	1.47 km to the north-east
Lime Kiln Hill Reservoirs	CWS	1.67 km north
Signal Hill Plantation Grassland	CWS	1.7 km north

Description

No description.

Contains more than 0.05 ha of CG3 upright brome calcareous grassland community. Supports frequent numbers of at least eight neutral grassland indicator species.

The site qualifies for habitat mosaic (over 10 ha in size supporting three or more habitat features), has an invertebrate index exceeding 500 and contains calcareous grassland supporting at least 0.05 ha of CG3 (upright brome) grassland.

This site qualifies as CWS because it supports a population of a Nationally Scarce plant species. Additionally, it supports type 10B waterbodies with beds of stoneworts *Charales* spp.

The site is listed as important in the bryophyte register for Cambridgeshire and has an overall invertebrate index exceeding 500.

Supports frequent numbers of at least 3 strong neutral grassland indicator species and frequent numbers of at least 6 strong calcareous grassland indicator species.

Supports populations of Nationally Scarce plant species (*Fumaria vaillantii*, *Fumaria parviflora*, *Linum perenne*). Also contains at least 0.05 ha of NVC CG3 grassland and frequent numbers of at least six strong calcareous grassland indicators species.

Site name	Designation	Approximate distance from Site	Description
Wort's Causeway Road Site Verge	CWS / Protected Road Verge	1.86 km to the north-west	Contains more than 0.05 ha of CG3 upright brome <i>Bromus erectus</i> calcareous grassland community. Supports frequent numbers of at least eight neutral grassland indicator species.
Cambridge University Botanic Garden	CWS	1.96 km north	The site is listed as a site of importance in the Bryophyte Site Register (NCC 1985) for Cambridgeshire and has an overall invertebrate index exceeding 500
Red Cross Lane Drain	City Wildlife Site (CiWS)	460 m to the east	Supports five or more neutral grassland indicator species in frequent numbers.
Hedgerow West of Babraham Road	CiWS	490m to the east	Hedgerow at least 100 m in length and 2 m in width at its widest point with four or more woody species.
Hobson's Brook South	CiWS	575 west	Chalk stream together with adjacent semi-natural habitat that has not been grossly modified through canalisation and/ or poor water quality.
Hobson's Brook Mid	CiWS	605 west	Chalk stream together with adjacent semi-natural habitat that has not been grossly modified through canalisation and/ or poor water quality.
Long Road Plantation	CiWS	1.08 km to the north	Recent woodland 1 ha or more in area and with five or more woodland plants.
Bentley Road Paddocks	CiWS	1.3 km north	Neutral grassland with two or more strong neutral grassland indicator species in frequent numbers.

Site name	Designation	Approximate distance from Site	Description
Hobson's Conduit / Vicar's Brook	CiWS	1.38 km north	Chalk stream together with adjacent semi-natural habitat that has not been grossly modified through canalisation and/ or poor water quality. Also supports a small group of pollard willows in a semi-natural setting
Clare Wood	CiWS	1.4 km north	Scrub over 0.5 ha in area with four or more woody species. Also qualifies as semi-natural habitat adjacent to a chalk stream.
Trumpington Road Woodland	CiWS	1.7 km to the north-west	Woodland 1 ha or more in area with five or more woodland plants.
Grantchester Road Plantations	CiWS	1.8 km to the north-west	Woodland over 1 ha in area and with five or more woodland plants.
Eight Acre Wood and Seven Acres Wood	CiWS	1.9 km west	Woodland over 1 ha in area and with 5 or more woodland plants.

Irreplaceable and priority habitats

10.10.63 The desk study found no ancient woodland/ veteran trees within the Site during the desk study. Native hedgerows and lowland mixed deciduous woodland are priority habitats within the RLB. Additionally, the River Granta was also identified as a priority river habitat.

Strategic significance

- 10.10.64 Part of the Site falls within the Gog Magog and Chalkland fringe; an area identified within the Greater Cambridge Local Plan First Proposal as being an area targeted for Policy BG/GI: Green infrastructure. This aims to provide links between habitats, connecting to the wider landscape.
- 10.10.65 Additionally, a B-line which is also noted as a pollinator corridor in the Greater Cambridge Local Plan First Proposal bisects the site between Sawston and Great Shelford. Therefore, habitats of rich pollen and nectar resource within this area including speciesrich grassland, species-rich hedgerows and mixed scrub have been given High strategic significance as they are formally identified in the local plan.
- 10.10.66 An area of lowland mixed deciduous woodland, identified as priority habitat from the desk study and located within Shelford - Haverhill disused railway CWS, west of High Street to the south west of Babraham lies within the RLB. This area of woodland connects to other similar habitat beyond the Site and as such has been given High strategic significance.

- 10.10.67 Cambridge City Council's Biodiversity Strategy 2022-2030 lists the following habitats that occur within the Site as a priority and are therefore given High strategic significance:
 - Lowland mixed deciduous woodland
 - Hedgerows
- 10.10.68 All other area habitats have been assessed as being of Medium with the exception of developed; sealed surface, vacant/ derelict land/ bare ground and productive land under the cropland broad habitat type, i.e. cereal crops winter stubble, cereal crops and noncereal crops which have been assigned Low strategic significance.

On-Site baseline condition assessment – Area habitat condition assessment

- 10.10.69 The majority of the Site comprised cropland composed of cereal and non-cereal crops. Other habitats include grassland comprising other neutral grassland and modified grassland predominantly in Good or Moderate condition, heathland and scrub comprising mixed scrub in Moderate condition, and a single non-priority pond in Poor condition. Other areas of the Site comprise existing roads and footpaths classified as developed land/ sealed surface for which there is no condition assessment. Urban trees are also present within the Site to the north predominantly classified as Poor owing to their immature age class.
- 10.10.70 Woodland is also present on Site comprising other broadleaved and lowland mixed deciduous woodland. An area of lowland mixed deciduous woodland was precautionarily assessed as Good condition as it was not accessed during surveys. Other areas of woodland were Poor or Moderate condition. A map of the habitat types provided in Appendix A.

On-Site baseline condition assessment – Linear hedgerow condition assessment

- 10.10.71 The Site comprises Native hedgerows predominately considered to be low distinctiveness in Moderate condition. Native hedgerow associated with bank or ditch and native hedgerow with trees of medium distinctiveness and Moderate or Good condition are also present. Native hedgerow associated with bank or ditch has been included as a hedgerow only in baseline unit calculations with the ditch assessed separately in the ditch assessment below.
- 10.10.72 Native species rich hedgerows present on Site predominantly comprised medium distinctiveness, without trees or banks/ ditches, in Moderate or Good condition. Native species rich hedgerow with trees of high distinctiveness in Good condition is also present.
- 10.10.73 Approximately 0.34 km of line of trees was recorded as being present on Site in either Poor condition due to their young age and being newly planted or as a precautionary Good condition as they were not accessed. A map of the hedgerow types is provided in Appendix B.

On-Site baseline condition assessment – River condition assessment

The river condition assessment found that all sub-reaches on the River Granta were of 10.10.74 River Type H (straight-sinuous planform with a sand channel bed). Detailed outputs of the river condition assessment are presented in Table A10.12.7.

Table A10.12.7 River type of the River Granta

River Type	Sub-reach								
indicators	D1-S1	D1-S2	D1-S3	D2-S1	D2-S2				
River category	Other	Other	Other	Other	Other				
Braiding index (A1)	1	1	1	1	1				
Sinuosity index (A2)	1.090	1.090	1.090	1.046	1.046				
Anabranching index (A3)	1	1	1	1	1				
Level of confinement (A4)	Unconfined	Unconfined	Unconfined	Unconfined	Unconfined				
Valley gradient (A5)	0.002	0.002	0.002	0.001	0.001				
Bedrock reaches (A6)	False	False	False	False	False				
Coarsest bed material size class (A7)	Cobble	Cobble	Cobble	Cobble	Cobble				
Average alluvial bed material size class (A8)	Sand	Sand	Sand	Sand	Sand				
Calculated River Type	Н	Н	Н	Н	Н				

10.10.75 The final River Granta condition scores for each sub-reach, that were inputted into the Natural England Biodiversity Metric 3.1, are listed in Table A10.12.8. Sub-reaches D1-S1 and D2-S2 were found to be Fairly Poor, with the other sub-reaches classed as Moderate (noting that the lower threshold for Moderate is 0.5, meaning that D1-S3 is borderline Fairly Poor/ Moderate).

Sub-reach	River type	Condition score	Condition class	River shape
D1-S1	Н	0.19	Fairly Poor	2.25
D1-S2	Н	1.40	Moderate	2.47
D1-S3	Н	0.53	Moderate	1.84
D2-S1	Н	0.91	Moderate	1.74
D2-S2	Н	0.41	Fairly Poor	1.66

Table A10.12.8 River condition scores and classes for the River Granta

- 10.10.76 In the UK, historical channel modifications mean that many river channels are 'overdeep' indicating that they are at least partially hydrologically and ecologically disconnected from their riparian margins and floodplains. These channels therefore have reduced potential to support biodiversity. Recognition of an overdeep channel is best assessed based on field investigation by an expert, but an index ('river shape' Table A10.12.8) has been created based on data collected during MoRPh surveys which can be used to assess the likelihood of a surveyed channel being sufficiently overdeep to affect its hydrological and ecological lateral connectivity. A river with a river shape score of ≤4 is likely to be overdeep, whilst rivers with a score of ≤2 are highly likely to be overdeep. All sub-reaches on the River Granta were therefore considered overdeep.
- 10.10.77 The Biodiversity Metric 3.1 Technical Supplement states that if a channel is considered to be overdeep, its river condition assessment should be reduced by one class, however, this guidance applies only where the condition is estimated to be Good or Fairly Good. Therefore, the condition class has not been downgraded for any of the River Granta sub-reaches. It should be noted, however, that the overdeep channel shape is likely to be reflected in the river condition indicator scores, and so a possible method for achieving biodiversity net gain may be to seek enhancements that address this.

On-Site baseline condition assessment – Ditch condition assessment

10.10.78 The ditch condition scores for each sub-reach are presented in Table A10.12.9. The condition of ditches was found to be predominantly Poor, but ditch W9 and Hobson's Brook (D4) were found to be at Moderate condition. W2 was also recorded as Moderate, but as the ditch was dry at the time of survey it was downgraded to Poor as this implies that sufficient water levels are not maintained (Criteria 6; "as a guide a minimum summer depth of approximately 50 cm in minor ditches").

Watercourse	Sub-reach	Number of criteria met	Condition class
Hobson's Brook	D4	6	Moderate
Unnamed ditches	D2E	3	Poor
	D5A	NA*	Poor
	W1	5	Poor
	W2	5	Poor
	W3	3	Poor
	W4	4	Poor
	W5	2	Poor
	W6	3	Poor
	W7	NA*	Poor
	W8	5	Poor
	W9	6	Moderate
	W22	NA*	Poor
	W24	4	Poor
	W27	NA*	Poor
	W29	4	Poor
*Condition class as	signed based on n	earby ditches aerial imagery	nast surveys and

Table A10.12.9 Ditch Condition Scores and Classes for Hobson's Brook and **Unnamed Ditches**

Condition class assigned based on nearby difches, aerial imagery, past surveys and professional judgement

10.10.79 Since Hobson's Brook needs to be considered under the rivers and streams broad habitat typology the ditch condition assessment does not apply. The condition class of Hobson's Brook has been assigned with reference to the completed ditch condition assessment and based on professional judgement as to how it would have been classified using the MoRPh survey method, in addition to review of:

- Photos of the Site
- Aerial imagery
- WFD status of the Hobson's Brook water body
- MoRPh results from the River Granta

10.10.80 As a result, an assumption of Fairly Poor condition has been applied to the Hobson's Brook, which aligns to the lowest classification recorded for the River Granta sub-reaches.

On-site baseline units – Area habitat baseline units

- 10.10.81 The area, distinctiveness, strategic significance and condition of area habitats were input to the Metric to calculate the Area Habitat Biodiversity Units (AHBU). The baseline conditions on Site currently generate a total of 278.23 AHBU over an area of approximately 114.17 ha.
- 10.10.82 A summary of the areas, condition, biodiversity units and retained habitat of the baseline area habitats are shown in Table A10.12.10. A UKHab plan of the baseline habitats is shown in Annex A.

Broad	Proposed habitat	Baseline	Distinctiveness		Condition		Strategic s	ignificance	Baseline units	Retained	Retained units
naditat		area (na)	Band	Score	Band	Score	Band	Score		area (na)	
Cropland	Arable field margins game bird mix	0.1791	Medium	4	N/A	1	Medium	1.10	0.79	0.0062	0.03
Cropland	Arable field margins pollen & nectar	1.9563	Medium	4	N/A	1	Medium	1.10	8.61	0	0
Cropland	Cereal crops winter stubble	18.3831	Low	2	N/A	1	Low	1.00	36.77	0.1304	0.26
Cropland	Cereal crops	57.2152	Low	2	N/A	1	Low	1.00	114.43	0.1407	0.28
Cropland	Non-cereal crops	21.8132	Low	2	N/A	1	Low	1.00	43.63	0	0
Grassland	Modified grassland	1.8570	Low	2	Good	3	Medium	1.10	12.26	0.0877	0.58
Grassland	Modified grassland	2.3532	Low	2	Moderate	2	Medium	1.10	10.35	0.0857	0.38
Grassland	Modified grassland	0.1852	Low	2	Poor	1	Medium	1.10	0.41	0.0116	0.03
Grassland	Other neutral grassland	0.2921	Medium	4	Good	3	Medium	1.10	3.86	0.0117	0.15
Grassland	Other neutral grassland	0.7695	Medium	4	Moderate	2	Medium	1.10	6.77	0.1254	1.10
Grassland	Other neutral grassland	0.1447	Medium	4	Moderate	2	High	1.15	1.33	0	0
Grassland	Other neutral grassland	0.0667	Medium	4	Poor	1	Medium	1.10	0.29	0	0
Grassland	Other neutral grassland	1.1606	Medium	4	Poor	1	High	1.15	5.34	0.0264	0.12
Heathland and shrub	Mixed scrub	0.0919	Medium	4	Moderate	2	High	1.15	0.85	0	0
Lakes	Ponds (Non- Priority Habitat)	0.0206	Medium	4	Poor	1	Medium	1.10	0.09	0.0150	0.07
Urban	Developed land; sealed surface	4.1450	V.Low	0	N/A - Other	0	Low	1.00	0	1.8462	0
Urban	Urban Tree	0.0056	Medium	4	Moderate	2	Medium	1.10	0.05	0.0056	0.05
Urban	Urban Tree	0.2842	Medium	4	Poor	1	Medium	1.10	1.25	0.2842	1.25
Urban	Vacant/derelict land/ bareground	0.0010	Low	2	Poor	1	Low	1.00	0	0	0
Woodland and forest	Lowland mixed deciduous woodland	0.6545	High	6	Good	3	High	1.15	13.55	0.6543	13.54

Table A10.12.10 Baseline area habitat scores and unit values within the Site

Broad	Proposed habitat	Baseline	Distinctiveness		Condition		Strategic significance		Baseline units	Retained	Retained units
Παριται		area (na)	Band	Score	Band	Score	Band	Score			
Woodland and forest	Lowland mixed deciduous woodland	0.0627	High	6	Moderate	2	High	1.15	0.87	0	0
Woodland and forest	Lowland mixed deciduous woodland	0.2502	High	6	Poor	1	High	1.15	1.73	0.1622	1.12
Woodland and forest	Other woodland; broadleaved	0.0003	Medium	4	Good	3	Medium	1.10	0	0	0
Woodland and forest	Other woodland; broadleaved	1.1320	Medium	4	Moderate	2	Medium	1.10	9.96	0.1645	1.45
Woodland and forest	Other woodland; broadleaved	1.1491	Medium	4	Poor	1	Medium	1.10	5.06	0.0111	0.05
Totals		114.17						278.23	3.77	20.46	

On-site baseline units – Linear hedgerow baseline units

- 10.10.83 The Site comprises a baseline of 73.53 hedgerow units over a length of 8.71 km.
- 10.10.84 A summary of the length, condition and biodiversity units and retained habitat of the baseline linear hedgerow are shown in Table A10.12.11.

Broad	Proposed habitat	Baseline	Distinctiveness		Condition		Strategic significance		Baseline	Retained	Retained units
nabitat		(km)	Band	Score	Band	Score	Band	Score	units	iength (na)	
Line of trees	Line of Trees	0.1009	Low	2	Good	3	Medium	1.10	0.67	0.1009	0.67
Line of trees	Line of Trees	0.2409	Low	2	Poor	1	Medium	1.10	0.53	0.2409	0.53
Hedgerow	Native Hedgerow (associated with ditch)	0.6679	Low	2	Good	3	High	1.15	4.61	0.5916	4.08
Hedgerow	Native Hedgerow (associated with ditch)	0.8112	Low	2	Moderate	2	High	1.15	3.73	0.7903	3.64
Hedgerow	Native Hedgerow with trees	0.7631	Medium	4	Good	3	High	1.15	10.53	0.6048	8.35
Hedgerow	Native Hedgerow with trees	0.6356	Medium	4	Moderate	2	High	1.15	5.85	0.6009	5.53
Hedgerow	Native Hedgerow	0.9663	Low	2	Good	3	High	1.15	6.67	0.8964	6.19
Hedgerow	Native Hedgerow	1.6514	Low	2	Moderate	2	High	1.15	7.60	1.5158	6.97
Hedgerow	Native Hedgerow	0.1943	Low	2	Poor	1	High	1.15	0.45	0.1506	0.35
Hedgerow	Native Species Rich Hedgerow	0.1284	Medium	4	Good	3	High	1.15	1.77	0.1284	1.77
Hedgerow	Native Species Rich Hedgerow with trees	0.6932	High	6	Good	3	High	1.15	14.35	0.2900	6.00
Hedgerow	Native Species Rich Hedgerow	0.7479	Medium	4	Good	3	High	1.15	10.32	0.7121	9.83
Hedgerow	Native Species Rich Hedgerow	0.3003	Medium	4	Moderate	2	High	1.15	2.76	0.2162	1.99
Hedgerow	Native Species Rich Hedgerow	0.8051	Medium	4	Poor	1	High	1.15	3.70	0.8051	3.70
Total		8.71							75.53	7.64	59.59

Table A10.12.11 Baseline linear hedgerow scores and unit values within the Site

On-site baseline units – Rivers and streams baseline units

- 10.10.86 The length, distinctiveness, strategic significance, encroachment and condition of the rivers and streams (River Granta and Hobson's Brook) were inputted into the Metric to calculate RBUs.
- 10.10.87 The River Granta has been identified as priority river habitat meaning that it is assigned a distinctiveness score of Very High. This is because the river is marked as a chalk river on Natural England's priority habitat chalk rivers map. Additionally, a review of the available aquatic ecological data and records of water-related mammals in the area found that five species from criterion level B (European eel, bullhead, brown trout, water vole, and otter) were present within the River Granta. There are also historical records of one criterion level A species (spined loach). Given that to qualify as priority habitat the watercourse needs to have records of any one criterion A species or any six criteria B species, the watercourse has been classified as priority river habitat. This is a precautionary approach which takes into account the high likelihood that spined loach are still a component of the watercourse's fish population.

- 10.10.88 The sub-reach of Hobson's Brook within the Site is assigned High distinctiveness based upon it not meeting any conditions for priority river habitat and therefore being entered into the Metric as Other Rivers and Streams. However, it should be noted that this sub-reach is a tributary of the main Hobson's Brook which is considered a chalk river.
- The River Granta has High strategic significance as it is a WFD reportable water body and 10.10.89 County Wildlife Site. This has been entered into the Metric as 'Within River Basin Management Plan'. Although Hobson's Brook is also a WFD water body, the watercourse at the point of interaction is well upstream of the delimited watercourse on the Environment Agency's Catchment Data Explorer. However, Hobson's Brook is still assigned High strategic significance as it is included in Local Plans due to its relation to Hobson's Brook City Wildlife Site and Nine Wells Local Nature Reserve, directly to the east of the RLB where the watercourse arises.
- 10.10.90 In-watercourse encroachment was assigned as no encroachment for all watercourse, excepting D1-S1 and D1-S3. These sub-reaches were assigned major in-watercourse encroachment on account of the presence of bed/ bank reinforcement at both sites, an intermediate deflector at D1-S1, and a major weir (i.e., a permanent impermeable impounding structure across the entire channel width) at D1-S3.

- 10.10.91 Most sub-reaches have no riparian encroachment (D1-S1, D2-S1 and D2-S2). However, unvegetated bankside footpaths alongside Hobson's Brook and D1-S3, mean that these sub-reaches have been assigned Moderate and Minor riparian encroachment respectively. The difference in encroachment severity reflects the different distances of the developments from the river.
- 10.10.92 River condition has been assigned to the River Granta sub-reaches based on the results of field surveys. For the Hobson's Brook, condition is based upon professional judgement and review of available data.
- 10.10.93 No off-site reaches have been assessed as part of this BNG assessment. There is 0.99 km of river within the RLB; 0.84 km of which is Very High distinctiveness habitat (River Granta), and the remainder of which (0.14 km) is High distinctiveness habitat (Hobson's Brook). The on-site baseline RBUs for rivers and streams are presented in Table A10.12.11. The total baseline RBUs for rivers and streams is 12.53 RB.

Watercourse	Location	Length km (within the RLB)	River type (Distinctiveness)	Strategic significance	Condition	Extent of watercourse encroachment	Extent of riparian encroachment	Baseline RBUs
River Granta (D1-S1)	Adjoining A11 and RLB at southern limit of Proposed Development	0.202	Priority Habitat (Very High)	Within River Basin Management Plan (High)	Fairly Poor	Major	No encroachment	1.39
River Granta (D1-S3)	Structure 8 crossing at Babraham	0.135	Priority Habitat (Very High)	Within River Basin Management Plan (High)	Moderate	Major	Minor	1.18
River Granta (D2-S1)	Immediately upstream of D2-S2, near the Structure 6 crossing	0.373	Priority Habitat (Very High)	Within River Basin Management Plan (High)	Moderate	No encroachment	No encroachment	6.86
River Granta (D2-S2)	Structure 6 crossing at Stapleford	0.132	Priority Habitat (Very High)	Within River Basin Management Plan (High)	Fairly Poor	No encroachment	No encroachment	1.82
Hobson's Brook (D4)	Structure 4 crossing near Addenbrooke's Road	0.144	Other Rivers and Streams (High)	Within Local Plans (High)	Fairly Poor	No encroachment	Moderate	1.27
Totals		0.986						12.53

Table A10.11.12 On-site baseline river biodiversity units for rivers and streams

On-site baseline units – Ditches on-Site baseline units

- 10.10.94 The length, distinctiveness, strategic significance, encroachment and condition of the ditches were inputted into the Metric to calculate RBUs.
- 10.10.95 The ditches were assigned Low distinctiveness, as the Biodiversity Metric considers all ditches to have Low distinctiveness. The ditches are also all considered to have Low strategic significance.

10.10.96 No ditches have any in-watercourse encroachment, but many of them have been assigned major riparian encroachment as they adjoin roads or railways. These ditches are: W1, W2, W7, W8, W9, W22, W24 and W29. All other ditches have no riparian encroachment.

10.10.97 Ditch condition has been assigned to ditches based on the results of the field survey. No off-site reaches have been assessed as part of this BNG assessment. Baseline RBUs were calculated for 15 ditches within the RLB, totalling 2.4 km of ditch. The on-site baseline RBUs for ditches are presented in Table A10.12.13. The total baseline RBUs for ditches is 9.04 RBUs.

Ditch	Location	Length km (within the Site)	River type (distinctiveness)	Strategic significance	Condition	Extent of watercourse encroachment	Extent of riparian encroachment	Baseline RBUs
D2E	Tributary of the River Granta near Structure 6	0.240	Ditches (Low)	Low	Poor	No encroachment	No encroachment	0.96
D5A	Near Graham's Road	0.159	Ditches (Low)	Low	Poor	No encroachment	No encroachment	0.64
W1	Near Francis Crick Avenue	0.011	Ditches (Low)	Low	Poor	No encroachment	Major	0.03
W2	Crossed by Structures 1, 2 and 3 near Addenbrooke's Road	0.153	Ditches (Low)	Low	Poor	No encroachment	Major	0.46
W3	Near Stapleford	0.121	Ditches (Low)	Low	Poor	No encroachment	No encroachment	0.48
W4	Crossed by Structure 7 at Sawston	0.174	Ditches (Low)	Low	Poor	No encroachment	No encroachment	0.70
W5	Near Sawston	0.472	Ditches (Low)	Low	Poor	No encroachment	No encroachment	1.89
W6	Crossed by Structure 5 at Stapleford, near River Granta	0.468	Ditches (Low)	Low	Poor	No encroachment	No encroachment	1.87
W7	Adjoining railway at Addenbrooke's Road	0.020	Ditches (Low)	Low	Poor	No encroachment	Major	0.06
W8	Near Addenbrooke's Road	0.153	Ditches (Low)	Low	Poor	No encroachment	Major	0.46
W9	Near Francis Crick Avenue	0.012	Ditches (Low)	Low	Moderate	No encroachment	Major	0.07
W22	Near Graham's Road	0.118	Ditches (Low)	Low	Poor	No encroachment	Major	0.35
W24	Near Francis Crick Avenue	0.099	Ditches (Low)	Low	Poor	No encroachment	Major	0.30
W27	Alongside Nine Wells LNR	0.175	Ditches (Low)	Low	Poor	No encroachment	No encroachment	0.70
W29	Near Francis Crick Avenue	0.024	Ditches (Low)	Low	Poor	No encroachment	Major	0.07
Totals		2.399						9.04

Table A10.12.13 On-site baseline river biodiversity units for ditches

On-Site impacted units – Area habitat on-Site impacted units

- 10.10.98 There is a total of 257.77 baseline AHBU lost on Site, predominantly cropland, modified grassland, other neutral grassland and other woodland; broadleaved. These habitats are being lost due to the creation of the new active transport route and associated infrastructure.
- 10.10.99 While cropland habitats including cereal, non-cereal and winter stubble is being lost to the Proposed Development this will be replaced when the compound and laydown areas are returned following construction.
- 10.10.100 Small areas of lowland mixed deciduous woodland will also be lost to the Proposed Development. This is predominantly marginal fringes adjacent to the compound areas. However, one section, north of Hinton Way and one section south of Holm Farm where the Proposed Development crosses the River Granta will be bisected by the Proposed Development. These areas connect to larger woodland beyond the Proposed Development boundary. Proposed planting includes the opportunity to create the same habitat adjacent to the woodland north of Hinton Way which will allow habitat creation to be aided by natural regeneration post construction while retained habitat along the River Granta will continue to provide ecological connectivity.
- 10.10.101 Non-priority ponds and mixed scrub will also be lost. Other units impacted on Site, including woodland and grassland will be lost to accommodate the Proposed Development.
- 10.10.102 Proposed habitat units may be impacted by additional management and pollution run-off from the Proposed Development during construction and operation. The likely effects have been taken into account in the condition of post-development AHBU.

On-Site impacted units – Linear hedgerow on-Site Impacted units

10.10.103 There are 58 hedgerows on Site within the RLB, totalling 8.36 km of hedgerow habitat plus a 0.34 km line of trees. The Proposed Development would result in the loss of approximately 0.54 km of native hedgerow and 0.52 km of native species-rich hedgerow. Losses are due to the construction of the Proposed Development and associated infrastructure and are largely based around existing road junctions and compound areas. Where hedgerows are lost there are currently no proposals to re-create hedgerows in existing locations. Instead, proposed hedgerows will largely border the proposed road and provide a barrier between the Proposed Development and adjacent land.

On-Site impacted units – Rivers and streams on-Site impacted units

- 10.10.104 The Proposed Development impacts relate to the construction of single span viaduct crossings over the River Granta and its floodplain at Structures 6 and 8 (corresponding to D2-S2 and D1-S3 sub-reaches respectively). There is also a single-span bridge crossing of the Hobson's Brook at Structure 4.
- 10.10.105 A desktop exercise was conducted to determine how the Proposed Development would impact the baseline indicator scores produced from the River Condition Assessment. This exercise took a 'worst-case' approach to ensure that impacts were not missed.
- 10.10.106 It was not possible to scenario model the impacts of the Proposed Development on the Hobson's Brook as the watercourse was not MoRPh surveyed. It is assumed that the impacts of the Proposed Development for this sub-reach would be similar to those at subreaches on the River Granta.
- 10.10.107 Impacts on river condition indicators can be summarised as follows:

- The primary impact of the new crossings would be an increase in shading from the • structures. Shading would reduce the local condition through loss of bank-top, bankface, and in-channel vegetation. It would also increase the extent of bare sediment on the banks:
- There would be a positive impact on the condition of the river through the shifting of riparian land use from arable agriculture to species rich grassland. The riparian habitat would no longer be managed for agriculture and would be restored to reflect a more natural habitat;
- It is assumed that there will be no in-channel works or requirements for bank protection or reprofiling. Therefore, no changes to artificial structure, bank profile or inchannel features have been included:
- The final location of piers (and therefore their distance from the bank-top) is to be confirmed in detailed design, but it is assumed that there would be no change to riparian encroachment for any sub-reach. The River Granta piers (Structures 6 and 8) would be positioned outside of the riparian zone (>10 m from the bank-top), meaning that there is no encroachment from the Proposed Development. The bridge piers at the Hobson's Brook crossing (Structure 4) are assumed to have moderate encroachment (i.e., be placed within 4 to 10 m of the bank-top) but this does not signify a change from the baseline, as the baseline watercourse encroachment is moderate due to the presence of a footpath on the right bank.
- 10.10.108 This desktop scenario modelling predicts that the Proposed Development would cause a reduction in condition score of 0.37 at D1-S3 and of 0.24 at D2-S2. This reduction corresponds to a reduction in condition class from Moderate to Fairly Poor for approximately 0.025 km of D1-S3 (as the baseline condition score was borderline Fairly Poor/ Moderate) but does not lead to a reduction in condition for D2-S2. Without river condition indicators for the Hobson's Brook it is not possible to determine the potential reduction in condition score at this sub-reach through a desktop exercise. However, based on the output of scenario modelling for D1-S3 and D2-S2, it has been assumed that there would be no reduction in condition class for this section of the Hobson's Brook due to the similarity of the designed modification at this site to that on the River Granta.
- 10.10.109 In Biodiversity Metric 3.1 it is not possible to record a reduction in condition for Rivers and Streams. It must be recorded as a loss of watercourse and creation of watercourse in poorer condition. However, this results in an over-estimation of RBUs that are lost due to a reduction in condition, because of the application of multipliers for creation, associated with time to target condition and the difficulty in creating habitat.
- 10.10.110 As part of this BNG assessment therefore, an additional scenario has also been undertaken where the penalties for time to target condition and difficulty in creating habitat have been removed from the calculation of RBUs loss caused by reduction in condition. This bespoke approach gives a more appropriate estimate of RBUs loss as, although there will be a reduction in condition, none of the river would be physically lost to the Proposed Development and so the time to target condition and difficulty of habitat creation do not apply as it is simply a change in condition applied to an existing reach. As there would be no loss of river channel, the Proposed Development is compliant with the rules of the Metric which state that losses of very high distinctiveness habitat cannot be adequately accounted for through the Metric.

On-Site impacted units – Ditches on-Site impacted units

10.10.111 There are 15 ditches within the RLB, totalling 2.4 km of ditch habitat. The total baseline RBUs for ditches is 9.04 RBUs.

10.10.112 The impacts of the Proposed Development on ditches can be summarised as follows:

- The Proposed Development would result in the loss of approximately 0.042 km and 0.005 km of W2 at Structures 1/2 and 3 respectively. 0.031 km of W4 at Structure 7. and 0.030 km of W6 at Structure 5 due to the culverting of these ditches underneath the embankments. This is accounted for within the Metric by counting these ditch lengths as 'loss' and then having 'creation' of new culverts¹⁰. This comes to a total of 0.108 km of ditch which would be culverted:
- There would be the loss of an existing ditch (W8) which currently adjoins Addenbrooke's Road. This ditch would be partially retained or recreated as a swale just outside of the footprint of the embankment, although whether the swale would meet the definition of a ditch according to BNG, and the length that would be retained/ recreated is to be confirmed in detailed design. For the purposes of BNG assessment, it has been assumed that the whole ditch length (0.153 km) would be lost;
- One ditch (W5) would gain major riparian encroachment as the CSET Proposed Development alignment would pass parallel to the existing ditch. To account for the riparian encroachment of the W5 ditch in the Metric, the ditch length has been entered as loss and has then been created with major riparian encroachment. There is no change in terms of encroachment for all other ditches.

On-Site post development – Area habitat

- 10.10.113 The new proposed public transport route, including the path for active travel, and travel hub facility providing car parking and bike storage are categorised as built linear features. Areas of modified grassland or other neutral grassland border the hub facility and public transport route. Lowland calcareous grassland planting is proposed in the central part of the Proposed Development where the public transport route bisects an exposed cutting.
- 10.10.114 Woodland creation is present around the hub facility acting as a landscape feature and create visual screening and connectivity between existing habitats.
- 10.10.115 The creation of ditches (see paragraph 10.10.134) has been incorporated into the landscape design for the Proposed Development. These will comprise a top width of 6 m. In-line with the Biodiversity Metric 3.1 User Guide, river and stream habitats greater than 5 m can be recorded as areas as well as lengths. If doing so, the area the watercourse occupies should be excluded from the ABU calculation. However, due to the ephemeral nature of ditches, it is anticipated that the mean water height within the ditches incorporated into the landscape design for the Proposed Scheme will not achieve this width for most of the year. Furthermore, the design of the ditches incorporates a stepped edge resulting in a narrow width from bank to bank when water levels are lower. Therefore, the ditches have been included as linear features only and not incorporated into the post development area habitat.
- 10.10.116 The post-work Site habitat creation is currently predicted to generate a total of 440.08 area habitat biodiversity units (AHBU) over an area of 110.4 ha.
- 10.10.117 Due to the duration of time for which compound areas will be occupied these habitats are considered to be lost and will be re-established once construction of the Proposed Development is complete. These areas predominantly comprise cropland.
- 10.10.118 This equates to a net gain of 182.31 AHBU and a percentage increase of 65.53%.

10.10.119 The overall unit and area change for the Site is shown in Table A10.12.14 with summary of post development habitat creation broken down by broad habitat type provided in Table A10.12.15. A plan showing the post-works habitat creation is shown in Annex B

¹⁰ Any river or stream that is culverted and present within the Order limits has a River Type of 'Culvert' and is automatically assigned a condition of 'Poor'.

Table A10.12.14 Summary of area results

Measurement	Habitat units	Area (ha)
Baseline value	278.23	114.17
Retained	20.46	3.77
Habitat creation	440.08	110.40

Total post construction	460.54	114.17
Change in biodiversity units (% change)	182.31 (65.53 %)	-

Table A10.12.15 On Site area habitat creation

Broad habitat	Proposed habitat	Proposed	Distinctivene	ss	Condition		Strategic significance		Habitat units
		area (na)	Band	Score	Band	Score	Band	Score	aeliverea
Urban	Lowland calcareous grassland	0.2019	High	6	Moderate	2	High	1.15	0
Cropland	ropland Modified grassland		Low	2	Poor	1	Medium	1.10	56.23
Cropland	Other neutral grassland	16.0897	Medium	4	Moderate	2	High	1.15	11.67
Cropland	Other neutral grassland	24.1711	Medium	4	Moderate	2	Medium	1.10	13.14
Grassland	Mixed scrub	0.1618	Medium	4	Moderate	2	Medium	1.10	0.64
Grassland	Ponds (Non- Priority Habitat)	1.8673	Medium	4	Moderate	2	Medium	1.10	3.41
Grassland	Lowland mixed deciduous woodland	1.3295	High	6	Moderate	2	High	1.15	123.87
Grassland	Other woodland; broadleaved	5.0114	Medium	4	Moderate	2	Medium	1.10	178.00
Heathland and shrub	Arable field margins game bird mix	0.1800	Medium	4	N/A	1	Medium	1.10	1.19
Lakes	Arable field margins pollen & nectar	2.0300	Medium	4	N/A	1	Medium	1.10	14.77
Woodland and forest	Lowland calcareous grassland	0.2019	High	6	Moderate	2	High	1.15	1.94
Woodland and forest	Modified grassland	1.6059	Low	2	Poor	1	Medium	1.10	25.84
Cropland	Other neutral grassland	16.0897	Medium	4	Moderate	2	High	1.15	0.76
Cropland	Other neutral grassland	24.1711	Medium	4	Moderate	2	Medium	1.10	8.62
Total area (ha)	ea (ha) 110.40 Total units					440.08			

On-Site post development – Linear hedgerow

- 10.10.120 New hedgerow creation will be incorporated into the landscape design for the Proposed Development. This will comprise native species-rich hedgerow species such as common dogwood, hazel, common hawthorn, wayfaring tree, blackthorn, and dog rose.
- 10.10.121 Hedgerows will predominantly form boundary features between the public transport route the surrounding land including agricultural fields.
- 10.10.122 Hedgerows and lines of trees retained total 7.64 km and 59.59 hedgerow biodiversity units (HBU). Hedgerow creation consisting of native species-rich hedgerow which totals 11.9 km, or 91.62 HBU, resulting a total net unit change of 77.67 HBU.

10.10.123 The overall unit and length change for the Proposed Development is shown in

Table A10.12.16. A plan showing the post-works habitat creation is shown in Appendix B. The results of the post development habitat creation are summarised in Table A10.12.17.

Table A10.12.16 Summary of linear hedgerow results

Measurement	Hedgerow Biodiversity Units	Length (km)
Baseline value	73.53	8.71
Retained	59.59	7.64
Hedgerow creation	91.62	11.90
Total post construction	151.20	19.54
Change in biodiversity units (% change)	77.67 (105.62 %)	-

Table A10.12.17 On-Site Habitat Creation - Linear Hedgerow

Broad habitat	Proposed habitat	Total length (km)	Distinctivenes	Distinctiveness Condition			Strategic signi	ficance	Habitat units delivered
			Band	Score	Band	Score	Band	Multiplier	
Hedgerow	Native Species Rich Hedgerow	11.9	Medium	4	Moderate	2	Low	1	79.67

On-Site post development – Rivers and stream

- 10.10.124 For the Rivers and Streams Module of the Metric, the Proposed Development design would result in a reduction of RBUs from 12.53 RBUs to 12.36 RBUs for the River Granta and Hobson's Brook combined, which equates to a 1.34% net loss.
- 10.10.125 The net loss relates to the reduction in condition class of 0.025 km of the River Granta at the Structure 8 crossing (D1-S3). A desktop exercise showed that the shading effect of the bridge crossing would cause a reduction in condition class from Moderate to Fairly Poor. At the Structure 6 crossing of the River Granta, the reduction in condition score did not correspond to a reduction in condition class. It was not possible to scenario model the impact of the Proposed Development on Hobson's Brook as the watercourse was not MoRPh surveyed but has been assumed that the Proposed Development would result in no impact on condition class, based on results from scenario modelling of D1-S3 and D2-S2.

Table A10.12.18 Rivers and Streams Biodiversity Net Gain Post-works

10.10.126 A breakdown of the Rivers and Streams RBUs delivered within the RLB are presented in Table A10.12.18.

Watercourse	Sub- reach	Baseline Length (within site boundary) (km)	Length Lost (km)	Length Retained (km)	Length Enhanced (km)	Length Created (km)	Baseline Units	Retained Units	Units Lost	Baseline Units Enhanced	Units Generated	Total Post-works Units
River Granta	D1-S1	0.202	0	0.202	0	0	1.39	1.39	0	0	0	1.39

	D1-S3	0.135	0.025 (at Moderate)	0.110	0	0.025 (at Fairly Poor)	1.18	0.96	0.22	0	0.05	1.01
	D2-S1	0.373	0	0.373	0	0	6.86	6.86	0	0	0	6.86
	D2-S2	0.132	0	0.132	0	0	1.82	1.82	0	0	0	1.82
Hobson's Brook	D4	0.144	0	0.144	0	0	1.27	1.27	0	0	0	1.27
Total (Rivers and Streams)		0.986	0.025	0.961	0	0.025	12.53	12.31	0.22	0	0.05	12.36 (-1.34%)
							•		•			-

Note: the reported numbers are as shown in the Metric.

- 10.10.127 Given the limitations with the Metric, which assumes that River Granta RBUs are lost to the Proposed Development and then created in poorer condition (rather than allowing for a reduction in condition without any loss of river), the loss in RBUs has been calculated without the multipliers for time to target condition and difficulty in creating habitat (see Section 3.4.3). This is likely to be more representative of the actual RBUs lost as part of the Proposed Development design as the watercourse would not be physically lost. This means that the post-works RBUs would be 12.47 (instead of 12.36 according to the Metric), meaning that there would be a 0.44% net loss. As this is a bespoke approach, no BNG calculator line entries can be provided that would capture this. It is included to demonstrate that the reported net loss of 1.34% is likely to be an overestimate of the net loss for Rivers and Streams.
- 10.10.128 There is an opportunity for enhancements to the River Granta and Hobson's Brook at subreaches which are within the RLB but are not affected by the crossings. By examining the river condition indicator scores of the River Granta sub-reaches, potential opportunities for enhancements have been identified specific to the Proposed Development. These are detailed in Table A10.12.19.

Table A10.12.19 Opportunities for enhancements in the River Granta and Hobson's Brook

Opportunity	Description and mechanism of improvement	Associated baseline indicators			
Removal of artificial	There are some small areas of bank reinforcement, a deflector	C7: Bank face artificial bank profile extent;			
structures	and a weir near the Babraham Crossing. There may be an	C8: Bank face reinforcement extent;			
	structures which are no longer required which would improve	C9: Bank face reinforcement material severity;			
	bank/bed complexity and flow naturalness.	E8: Channel bed reinforcement extent;			
		E9: Channel bed reinforcement materials severity;			
		E10: Channel bed artificial features severity.			
Bank reprofiling	The River Granta is incised with uniform bank profiles,	C1: Bank face riparian vegetation structure;			
	therefore banks could be reprofiled with the aim of creating a more natural two-	C3: Bank face natural bank profile extent;			
	stage channel to facilitate the establishment of marginal	C4: Bank face natural bank profile richness;			
	features (such as berms/ benches). This could also	C7: Bank face artificial bank profile extent;			
	between the river and its floodplain and could benefit	D1: Channel margin aquatic vegetation extent;			
	marginal and vegetation.	D3: Chanel margin physical feature extent;			
		D4: Channel margin physical feature richness.			

Opportunity	Description and mechanism of improvement	Associated baseline indicators
Introduction of woody material	Any suitable trees which are felled as part of the Proposed Development could be placed on the banks and within the channel to create flow deflectors. This would create fish habitat and contribute towards an increase in flow variability.	E3: Channel bed hydraulic feature richness; E4: Channel bed natural physical features extent.
Creation of in- channel geomorphological features	Riffle and pool features could be created by introducing coarse substrates (which would also increase substrate diversity) and wood features could be introduced as previously described to create and maintain pool features through scour.	E3: Channel bed hydraulic feature richness.E4: Channel bed natural physical features extent;E6: Channel bed material richness.

- 10.10.129 Scenario modelling has been undertaken to stress test the potential for watercourse enhancements within the RLB to provide net gain by increasing the number of post-works RBUs. This found that works could increase the condition of all sub-reaches which are not directly affected by the Proposed Development by one condition class (see Volume 3, Appendix 10.13 for stress test details).
- 10.10.130 The enhancements would need to occur alongside the removal of artificial structures, which are the existing cause of major watercourse encroachment, in a least one of the two sub-reaches in which they occur (D1-S1 and D1-S3). It should be noted that there is currently a high level of uncertainty around the feasibility of the required structure removal. If achievable there is the potential for watercourse enhancements, including structure removal, to yield at least a 20% net gain for the rivers and streams RBU component of the Metric.
- 10.10.131 Enhancements have not been included in the BNG assessment as it is not part of the current design, but these or similar enhancements could be incorporated into future design changes in order to meet Greater Cambridge Partnerships' BNG targets.

On-Site post development – Ditches

- 10.10.132 The Proposed Development would result in a net loss of RBUs relating to ditches from 9.04 to 7.83 RBUs, which equates to a 13.37% loss in RBUs.
- 10.10.133 This is a result of the culverting of three ditches (approximately 0.108 km in total), which is represented in the Metric as loss of ditch and creation of culvert. RBUs are also lost due to the loss of 0.153 km of W8, and the major riparian encroachment of W5 (approximately 0.472 km), which is represented in the Metric as loss of ditch length and then creation of the ditch with major encroachment.
- 10.10.134 Given site constraints and the hydrology of the existing ditches, there is limited potential for any enhancements to existing ditches to improve their condition class. However, new ditch creation will be incorporated into the landscape design for the Proposed Development. This will comprise of 6 m wide (top width) ditches designed to maintain sufficient depth throughout the year, with marginal berms incorporated into the design in order to support a range of emergent, submerged and floating-leaved plants.
- 10.10.135 A matrix of ditches has been incorporated into the landscape design at two locations:
 - Approximately 170 m length of ditches connecting wildlife ponds south of the Travel Hub; and
 - Approximately 489 m of ditches adjoining the River Granta (Stapleford) Crossing.
- 10.10.136 It is assumed that through the design principles being applied that created ditches would be able to achieve Moderate condition. This is a realistic assumption given that all eight ditch condition criteria must be met to achieve Good condition (Paragraph 10.10.21). Achieving all eight criteria may not be possible at this site given the presence of INNS within the catchment and the lowland, eutrophic nature of the adjacent river.
- 10.10.137 For this assessment, it is assumed that ditch creation will be delayed by two years as a worst case scenario (the duration of the anticipated construction programme) from the time of the initial Site clearance and the standard time to target condition has been adjusted accordingly. No delay in starting habitat creation has been applied to impacted units.
- 10.10.138 The retained ditches total 1.029 km in length and 3.88 river biodiversity units (RBU). Ditch creation of Moderate condition ditch totals 0.659 km, or 4.11 RBU, resulting a total net unit change of plus 2.90 RBU, or a 32.06% net gain.
- 10.10.139 A breakdown of the RBUs including the overall unit and length change for the Proposed Development is presented in Table A10.12.20. A plan showing the post-works habitat creation is provided in Annex B.

Watercourse	Baseline Length (within site boundary) (km)	Length Lost (km)	Length Retained (km)	Length Enhanced (km)	Length Created (km)	Baseline Units	Retained Units	Units Lost	Baseline Units Enhanced	Units Generated	Total Post- works Units
D2E	0.240	0	0.240	0	0	0.96	0.96	0	0	0	0.96
D5A	0.159	0	0.159	0	0	0.64	0.64	0	0	0	0.64
W1	0.011	0	0.011	0	0	0.03	0.03	0	0	0	0.03

Table A10.12.20 Ditches biodiversity net gain post-works

Watercourse	Baseline Length (within site boundary) (km)	Length Lost (km)	Length Retained (km)	Length Enhanced (km)	Length Created (km)	Baseline Units	Retained Units	Units Lost	Baseline Units Enhanced	Units Generated	Total Post- works Units
W2	0.153	0.047	0.106	0	0.047	0.46	0.32	0.14	0	0.07	0.39
W3	0.121	0	0.121	0	0	0.48	0.48	0	0	0	0.48
W4	0.174	0.031	0.143	0	0.031	0.70	0.57	0.12	0	0.04	0.62
W5	0.472	0.472	0	0	0.472	1.89	0	1.89	0	1.37	1.37
W6	0.468	0.030	0.438	0	0.030	1.87	1.75	0.12	0	0.04	1.80
W7	0.020	0	0.020	0	0	0.06	0.06	0	0	0	0.06
W8	0.153	0.153	0	0	0	0.46	0	0.46	0	0	0
W9	0.012	0	0.012	0	0	0.07	0.07	0	0	0	0.07
W22	0.118	0	0.118	0	0	0.35	0.35	0	0	0	0.35
W24	0.099	0	0.099	0	0	0.30	0.30	0	0	0	0.30
W27	0.175	0	0.175	0	0	0.70	0.70	0	0	0	0.70
W29	0.024	0	0.024	0	0	0.07	0.07	0	0	0	0.07
Travel hub ditches	0	0	0	0	0.170	0	0	0	0	1.06	1.06
River Granta (Stapleford) Crossing ditches	0	0	0	0	0.489	0	0	0	0	3.05	3.05
Total (Ditches)	0.240	0.733	1.666	0	1.239	9.04	6.31	2.73	0	5.63	11.94 (32.06%)
Note: the repor Note: Minor dis	ted numbers are as s crepancies in the add	hown in the Metr	ric. ction of RBUs are	e a result of rour	nding to two decir	mal places.					

10.10.140 If post-works RBUs are calculated without the multipliers for time to target condition and difficulty in creating habitat for W5 (as the only impact is an increase in riparian encroachment), this results in an increase of post-works RBUs of 11.99 compared to 11.94 RBUs, meaning that there would be 32.59% net gain instead of 32.06% according to the Metric. As ditches are low difficulty habitats to create according to the Metric, there is no penalty for difficulty in creating habitat applied, and so there is little difference in the calculated post-works RBUs for ditches between this bespoke approach and the results using the Metric.

10.10.141

10.10.142 Note that a conservative approach has been taken when calculating the length of ditch creation required to achieve the BNG target. A precautionary approach was taken on whether to include ditches which were dry at the time of survey in the calculation of RBUs, which may have led to an overestimation of baseline RBUs. Additional survey data would be needed to establish whether or not this assumption is appropriate.

On-Site post development – Strategic significance

- 10.10.143 Cambridge City Council's Biodiversity Strategy 2022-203019 lists lowland calcareous grassland as a priority. This habitat is proposed within the post-development landscape and planting plan and are therefore given High strategic significance.
- 10.10.144 Likewise lowland mixed deciduous woodland and hedgerows throughout the Site have been given High strategic significance and habitats containing pollen and nectar rich resources including species-rich grassland, species-rich hedgerows and mixed scrub within the Gog Magog and Chalkland Fringe area, pollinator corridor and B-Line have been given High strategic significance in line with baseline strategic significance.
- 10.10.145 All other natural habitats which fall under the broad habitats woodland and forest, lakes and grassland have been given Medium strategic significance while developed land; sealed surface has been given Low.

Trading summary

- 10.10.146 Biodiversity Metric 3.1 sets out a number of principles and rules that should be followed when undertaking BNG assessments. Rule 3 states:
- 10.10.147 'Trading down' must be avoided. Losses of habitat are to be compensated for on a "like for like" or "like for better" basis. New or restored habitats should aim to achieve a higher distinctiveness and/or condition than those lost.
- 10.10.148 Losses of irreplaceable or very high distinctiveness habitat cannot adequately be accounted for through the metric.
- 10.10.149 Biodiversity Metric 3.1 provides indicative advice regarding the actions to be taken to address habitat loss.

Trading summary – Very high distinctiveness habitats

- 10.10.150 For very high distinctiveness habitats losses cannot be adequately accounted for through the metric. For very high distinctiveness habitats where losses cannot be avoided, it is recommended that bespoke compensation is put in place.
- 10.10.151 The Proposed Development is predicted to result in a unit change of -0.17 units of priority river habitat. However, there would be no loss of direct habitat of very high distinctiveness, as the unit reduction is driven by a localised reduction in condition associated with viaduct crossing. This satisfies the trading rules.

Trading summary – High distinctiveness habitats

- 10.10.152 For high distinctiveness habitats, to avoid trading down Natural England recommends that habitat creation aims to create an equivalent number of units of the same habitat type.
- 10.10.153 The Proposed Development is predicted to result in a unit change of +0.44 units of lowland mixed deciduous woodland habitat and +0.57 units of calcareous grassland. There would be no change in units for river habitat of high distinctiveness. This satisfies the trading rules.

Trading summary – Medium distinctiveness habitats

10.10.154 For medium distinctiveness habitats, to avoid trading down Natural England recommends that habitat creation aims to create an equivalent number of units of the same broad habitat type. For example, bramble scrub can be replaced with mixed scrub without risking trading down.

10.10.155 The Proposed Development results in a total of +313.08 Medium distinctiveness area habitat units, this comprises +0.02 units of cropland, +285.66 units of grassland, +0.35 units of heathland and scrub, +14.74 units of lakes and +12.32 units of woodland and forest. This satisfies the trading rules for Medium distinctiveness area habitat.

Trading summary - Low distinctiveness habitats

- 10.10.156 For low distinctiveness habitat, to avoid trading down the guidance within the Metric recommends that habitat creation aims to create an equivalent number of units for habitats of the same distinctiveness or better. There is no requirement for this to come from the same broad habitat type. Units can be used from any higher distinctiveness habitat (medium, high or very high), provided they have not already been counted as part of the trading down for these higher distinctiveness habitats.
- 10.10.157 The Proposed Development is predicted to result in the overall unit loss of -131.87 low distinctiveness habitat area units. This comprises cropland and modified grassland. However, there is a sufficient surplus in the Medium distinctiveness habitats to account for this and therefore there is no trading down of low distinctiveness habitats.
- 10.10.158 No further action is required to offset trading down in low distinctiveness habitats.
- 10.10.159 The Proposed Development also results in a gain of 32.06% low distinctiveness habitat units relating to ditches.

Conclusion

- 10.10.160 The Biodiversity Metric 3.1 has been applied to the Proposed Development at the Site, based on the design at the time of publication of this report.
- 10.10.161 The assessment shows that the Proposed Development is predicted to result in a net gain of 182.31 AHBU, with a predicted total net change of +65.53%. The net change in hedgerow units is +77.67 which is a net gain of 105.62%. AHBU and HBU achieve and surpasses the 20% biodiversity net gain target for the Proposed Development. The Proposed Development is also predicted to result in a net gain of 11.94 RBUs (+32.06% gain) for ditches.
- 10.10.162 However, the assessment also shows that the Proposed Development is predicted to result in a net loss for the rivers and streams RBU component of the Metric, with a predicted total net change of -1.32%. This BNG assessment summarises enhancement opportunities that have been stress tested for their feasibility that could be applied to subreaches of the River Granta and Hobson's Brook (Volume 3, Appendix 10.13). To achieve the 20% net gain for the rivers and streams metric, existing encroachments such as weirs and other structures would need to be removed. These are partially outside of the Site boundary and their purpose is unknown so there is considerable uncertainty as to whether their removal would be possible. It is therefore unlikely that the 20% net gain for rivers and streams could be achieved and biodiversity credits would need to be purchased to offset biodiversity loss and achieve the 20% target.

- 10.10.163 The strategy for achieving BNG, including its definition in relation to the proposed biodiversity improvement parameters, will need to be discussed and agreed with local stakeholders and must be based on ecological functionality with regard to local conservation priorities and local biodiversity targets.
- 10.10.164 The undertaking of this BNG assessment has followed the mitigation hierarchy and the principles and rules of Biodiversity Metric 3.1.
- 10.10.165 A Landscape and Ecology Management Plan detailing appropriate BNG management and maintenance plan has been produced (ES Volume 3, Appendix 2.2). This details the measures required to ensure the habitat creation proposals reach their target condition and give confidence that every effort will be made to enable the Proposed Development to deliver the predicted net gain in biodiversity units.







		D	50 100 m	
tkins Limited @ 2 /	Atlantic Square, York Stree	et, Glaagow, G2 8JQ	Original Scale: 1:3,500	
ase Mapping: ©	OpenStreetMap and its	contributors	ATKINS	
rawn by: EM ate: 30/03/23	Checked by: NC Date: 30/03/23	Reviewed by: AW Date: 30/03/23	Member of the SNC Cavelin Group	





					×
tive I tive I	hedgeraw Hedgeraw with trees				
			D	50	100 m
	Atkins Limited @ 2 Atla	ntic Square, York Street,	Glasgow, G2 83Q	Original :	Scale: 1:3,500
	Base Mapping: @ Op	xenStreetMap and its o	ontributors	1	TKINS
	Drawn by: EM Date: 30/03/23	Checked by: NC Date: 30/03/23	Reviewed by: AW Date: 30/03/23	Mo	inder of the SNC Cavelin Group











ANNEX B SITE HABITAT CREATION









					Å
					~ `
	on- Priority Habitath				
•	idgerow				
			0	50 100 m	
	Atkins Limited @ 2 Atlantic	Square, Tork Street, (Glangow, G2 83Q	Original Scale:	1:3,500
	Base Mapping: @ OpenStree Drawn by: EM On	etMap and its contrib ecked by: NC	Reviewed by: AW	ΛΤΚΙ	NS
1				Plentar of the OLD on	









11			
			Å
			1 Martin
	1		
	3		
\sim	a states		
	where we have a set of some share	and the state	
		12.20	
	1.1.1.1.1	Y	1
		1	
	2.20		
	K		
10		7	August .
		I. Cam	
iority Habitat)			
w			
			50 100 m
Atkins Limited @ 2 Atla	ntic Square, Tork Street,	Glangow, G2 83Q	Original Scale: 1:3,500
Base Mapping: © Open	StreetMap and its contrib	utoni	ATKING
Drawn by: EM Date: 01/06/23	Checked by: NC Date: 12/09/23	Reviewed by: AW Date: 12/09/23	Perior of the OCCurrent Const



ANNEX C SUB-REACH RIVER CONDITION INDICATOR SCORES FROM MORPH SURVEYS OF THE RIVER GRANTA

Reach name:			D1-S1	D1-S2	D1-S3	D2-S1	D2-S2
	Vegetation structure	B1	2	2	2	2	2
d	Tree feature richness	B2	0	0	0	0	0
ank to	Water related features	B3	0	0	0	0	0
B	Invasive species	B4	-1	0	0	0	0
	Managed ground cover	B5	-3	-2	-2	-3	-2
	Riparian vegetation structure	C1	1	2	2	2	2
	Tree feature richness	C2	1	2	2	1	1
	Natural bank profile extent	C3	2	2	2	1	0
	Natural bank profile richness	C4	2	3	3	2	0
nk face	Natural bank material richness	C5	3	1	1	1	1
Ba	Bare sediment extent	C6	4	4	4	3	1
	Artificial bank profile extent	C7	-2	0	-2	-2	-2
	Reinforcement extent	C8	-2	0	-2	0	0
	Reinforcement material severity	C9	-1	0	-2	0	0
	Invasive species	C10	-1	-1	0	-2	-2
	Aquatic vegetation extent	D1	2	2	2	2	2
r margin	Aquatic morphotype richness	D2	2	2	2	2	2
Wate	Physical feature extent	D3	2	3	2	1	1
	Physical feature richness	D4	2	2	1	1	1

Reach name:			D1-S1	D1-S2	D1-S3	D2-S1	D2-S2
	Artificial features	D5	-2	0	-1	0	0
	Aquatic morphotype richness	E1	1	0	2	3	3
	Tree feature richness	E2	2	2	2	1	1
	Hydraulic richness	E3	1	1	2	1	0
	Natural feature extent	E4	0	0	0	2	0
_	Natural feature richness	E5	0	0	0	1	0
anne	Material richness	E6	3	3	0	3	1
Ch	Bed siltation	E7	-2	0	0	0	0
	Reinforcement extent	E8	-1	0	0	0	0
	Reinforcement severity	E9	-2	0	0	0	0
	Artificial feature severity	E10	0	0	-4	0	0
	Invasives	E11	-1	0	0	-1	0
	Filamentous algae E12		0	0	0	0	-1
	River Shape		2.25	2.47	1.84	1.74	1.66
	Average width (m)		5.0	4.6	5.0	4.6	5.0
	Average positive index		1.58	1.63	1.53	1.53	0.95
otals	Average negative index		-1.38	-0.23	-1	-0.62	-0.54
F	Condition Score		0.19	1.4	0.53	0.91	0.41
	River Type		Н	Н	Н	Н	Н
	Final Condition Class		Fairly Poor	Moderate	Moderate	Moderate	Fairly Poor

ANNEX D RESULTS OF DESKTOP EXERCISE TO ESTIMATE WORST CASE IMPACTS OF THE PROPOSED DEVELOPMENT ON THE RIVER GRANTA

Sub-reach name:			BASELINE	SCENARIO	BASELINE	SCENARIO
		D2-S2	D2-S2 crossing	D1-S3	D1-S3 crossing	
	Vegetation structure	B1	2	0	2	0
d	Tree feature richness	B2	0	0	0	0
ank to	Water related features	B3	0	0	0	0
ä	Invasive species	B4	0	0	0	0
	Managed ground cover	B5	-2	0	-2	0
	Riparian vegetation structure	C1	2	0	2	0
	Tree feature richness	C2	1	1	2	2
	Natural bank profile extent	C3	0	0	2	2
	Natural bank profile richness	C4	0	0	3	3
nk face	Natural bank material richness	C5	1	1	1	1
Ba	Bare sediment extent	C6	1	3	4	4
	Artificial bank profile extent	C7	-2	-2	-2	-2
	Reinforcement extent	C8	0	0	-2	-2
	Reinforcement material severity	C9	0	0	-2	-2
	Invasive species	C10	-2	-2	0	0
gin	Aquatic vegetation extent	D1	2	0	2	0
iter man	Aquatic morphotype richness	D2	2	0	2	0
Wa	Physical feature extent	D3	1	1	2	2

Sub-reach name:			BASELINE	SCENARIO	BASELINE	SCENARIO
			D2-S2	D2-S2 crossing	D1-S3	D1-S3 crossing
	Physical feature richness	D4	1	1	1	1
	Artificial features	D5	0	0	-1	-1
	Aquatic morphotype richness	E1	3	0	2	0
	Tree feature richness	E2	1	1	2	2
	Hydraulic richness	E3	0	0	2	2
	Natural feature extent	E4	0	0	0	0
	Natural feature richness	E5	0	0	0	0
annel	Material richness	E6	1	1	0	0
Ch	Bed siltation	E7	0	0	0	0
	Reinforcement extent	E8	0	0	0	0
	Reinforcement severity	E9	0	0	0	0
	Artificial feature severity	E10	0	0	-4	-4
	Invasives	E11	0	0	0	0
	Filamentous algae	E12	-1	0	0	0
	Average positive index		0.95	0.47	1.53	1.00
	Average negative index		-0.54	-0.31	-1.00	-0.85
otals	Condition Score		0.41	0.17	0.53	0.15
Τ	River Type		Н	Н	Н	Н
	Final Condition Class		Fairly Poor	Fairly Poor	Moderate	Fairly Poor